

12

EUROPEAN PATENT SPECIFICATION

45 Date of publication of the patent specification:
07.11.90

51 Int. Cl. ⁸: **A61K 31/485**

21 Application number: 88311290.6

22 Date of filing: 29.11.88

54 **Pharmaceutical compositions.**

30 Priority: 03.12.87 GB 8728294

43 Date of publication of application:
07.06.89 Bulletin 89/23

45 Publication of the grant of the patent:
07.11.90 Bulletin 90/45

84 Designated Contracting States:
AT BE CH DE FR IT LI LU NL SE

56 References cited:
EP-A-0 185 472

73 Proprietor: **RECKITT & COLMAN PRODUCTS LIMITED**,
One Burlington Lane, London W4 2RW(GB)

72 Inventor: **Lewis, John William**, 20 Sidings Court Skillings
Lane, Brough North Humberside(GB)

74 Representative: **Hardisty, David Robert et al, BOULT,**
WADE & TENNANT 27 Fumival Street, London EC4A
IPQ(GB)

EP 0 319 243 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

Description

This invention relates to compositions useful for the treatment of opiate dependence and more particularly to compositions containing naltrexone and buprenorphine.

5 Naltrexone (INN for 1-N-cyclopropylmethyl-14-hydroxynordihydromorphinone) is a pure opiate antagonist which, when administered orally (50 mg/day) as a maintenance drug for opiate addicts, blocks the effects of self-administered opiates and this contributes to the extinction of drug craving. Unfortunately, only about 10 per cent of addicts inducted on to a naltrexone treatment regime remain in treatment since naltrexone has no positive reinforcing effect to satisfy the needs of the addict. It also has the disadvantage that it precipitates abstinence in opiate abusers including those with only a low level of physical dependence. Thus an addict must be detoxified and be drug free for at least ten days before starting naltrexone treatment.

10 Buprenorphine (INN for N-cyclopropylmethyl-7&-[1-(S)-hydroxy-1,2,2-trimethylpropyl]6,14-endoethano-6,7,8,14-tetrahydronoripavine) has been shown in man to be a potent antagonist analgesic lacking the psychotomimetic effects found with other antagonist analgesics. Buprenorphine effectively relieves moderate to severe pain in doses of 0.1 mg or more administered either parenterally or sublingually.

The optimum therapeutic range for single doses is 0.3 mg - 0.6 mg by injection and 0.1 mg - 0.4 mg for sublingual tablets. In animal tests and in man buprenorphine has been shown to have both agonist (morphine-like) and (morphine) antagonist properties. However from direct dependence studies in animals and in man it has been concluded that buprenorphine does not produce significant physical dependence and the potential to produce psychological dependence is low as indicated by animal self administration studies and by the measurement of euphorogenic effects in human post addicts. In man the agonist and narcotic antagonist characteristics of buprenorphine have been demonstrated in opiate addicts. Thus oral buprenorphine in the dose range 6-16 mg has been shown to precipitate abstinence in highly dependent opiate addicts presenting for detoxification. On the other hand in a study involving subjects stabilised on a relatively low daily dose of oral methadone, sublingual buprenorphine could be substituted for methadone with only a low level of discomfort. In this situation buprenorphine was behaving as an opiate agonist of low intrinsic activity.

Thus buprenorphine has many of the desired characteristics of a treatment for opiate dependence (a) the ability to substitute for opiates in moderately dependent individuals (b) provide limited, long-lasting reinforcing (euphorogenic) effects which are acceptable to addicts (c) produce very mild abstinence effects when the drug is withdrawn, and (d) provide very good safety. With respect to (c) and (d) buprenorphine is markedly superior to methadone which is the only opioid agonist presently used for maintenance therapy.

35 For maintenance treatment there is the need for a product which can be safely administered on a "take home" basis. One of the potential problems of a sublingual buprenorphine product for the treatment of opiate addicts is its vulnerability to diversion if it is made available as a "take home" medication. Since the sublingual preparations to be absorbed have to be totally and relatively easily soluble, an addict in treatment could dissolve up the product and inject it. The useful sublingual dose range of buprenorphine for addict treatment (2 mg - 8 mg) is about ten times higher than the analgesic dose range and when injected is potentially equivalent to 60-240 mg morphine or 30-120 mg heroin; as such it will have a significant value to street addicts. It is therefore to be expected that if such a sublingual buprenorphine product were made available as "take home" medication a proportion of it would fall into the hands of street opiate users. Injection of the diverted buprenorphine would negate the primary purpose of a treatment for opiate dependence - to prevent intravenous drug use which is a major source of AIDS infection.

45 Our US Patent No 4661492 describes and claims in particular a method of treating pain which comprises the administration to a patient of a sublingually effective unit dosage of buprenorphine wherein the weight of buprenorphine is between 0.1 to 0.4 mg and simultaneously an amount of naltrexone sufficient to precipitate abstinence and thus prevent substitution in an opiate dependent subject, the weights of naltrexone and buprenorphine administered sublingually being within the ratio of 1:4 to 1:2. There is also disclosed and claimed an analgesic composition in sublingual unit dosage form comprising an active dose of buprenorphine of from about 0.1 to about 0.4 mg and an amount of naltrexone sufficient to prove aversive to a narcotic addict by parenteral administration but insufficient to compromise the analgesic action of the buprenorphine, the weight of naltrexone and buprenorphine being within the ratio of 1:4 to 1:2. The analgesic effect of these combination doses was equal to that of the equivalent dose of buprenorphine alone; however the ability of the combinations to precipitate abstinence in opiate-dependent subjects when injected was as great as that of the equivalent doses of naltrexone. Consequently an opiate dependent abuser would be discouraged from injecting the combination.

55 We have now found that there are doses of buprenorphine and naltrexone which when co-administered may be used in the treatment of opiate addicts.

60 According to this invention there is provided a pharmaceutical composition in sublingual unit dosage form for maintenance treatment of opiate addicts comprising from 2 to 8 mg buprenorphine and an amount of naltrexone sufficient to substantially attenuate the euphorogenic effect of the buprenorphine when injected and to provide greater opiate blocking effect than that of naltrexone alone wherein the weights of naltrexone and buprenorphine are the ratio of 1:4 to 1:1 the weights of buprenorphine and naltrexone be-

ing expressed in terms of weights of the base forms. The preferred weight of buprenorphine is 4 mg and that of naltrexone is in the range of 1 to 4 mg.

In an aspect of the invention there is provided a pharmaceutical composition in sublingual unit dosage form for maintenance treatment of opiate addicts comprising from 2 to 8 mg buprenorphine and an amount of naltrexone sufficient to substantially attenuate the euphorogenic effect of the buprenorphine when injected and to provide greater opiate blocking effect than that of naltrexone alone wherein the amount of naltrexone is in the range of 2 to 8 mg, the weights of buprenorphine and naltrexone being expressed in terms of weights of the base forms. The preferred weight of buprenorphine is 4 mg and that of naltrexone is in the range of 2 to 4 mg.

It is to be understood that the use of the terms buprenorphine and naltrexone comprehend not only the bases but also their pharmaceutically acceptable salts. Particular preferred salts are the hydrochlorides.

The sublingual combinations of the present invention contain in unit doses greater amounts of naltrexone than the sublingual analgesic compositions of our earlier invention. This amount of naltrexone exerts a substantial opiate antagonist effect sublingually as well as when injected and in this respect the drug abuse treatment combinations differ from the analgesic combinations. The addiction treatment combinations will have substantially less reinforcing (euphorogenic) effect than buprenorphine alone particularly when injected and will not be attractive to any opiate abuser even those who are not physically dependent. They will also act sublingually in precipitating abstinence in dependent individuals. For this reason it is desirable that opiate addicts should first be stabilised on buprenorphine alone before transfer to the combination. This transfer can be made without a drug free period as is necessary in the present procedure for transferring addicts from opiates to oral naltrexone since there is no precipitated abstinence when buprenorphine-maintained subjects are treated with naltrexone.

The preparations of the present invention are superior to equivalent preparations of buprenorphine alone for maintenance treatment of opiate dependence since they can be safely dispensed for "take home" use without fear of diversion. Furthermore we have shown that the abstinence effects following repeated administration of the combination are of an even lower level than those associated with buprenorphine alone.

The preparations of the present invention are superior to the present oral naltrexone maintenance product

(a) based on their limited level of agonist effect which makes them more acceptable to addicts and therefore gives improved rates of retention in treatment.

(b) based on our finding that they have greater ability to block the acute effects of opiates. This is a most important factor determining the efficacy of a maintenance treatment for opiate dependence.

In the rat tail pressure test (Green and Young, Br. J. Pharmac., 6, p572, 1957), dose-response curves for morphine were determined after four days' pretreatment with twice-daily subcutaneous doses of saline, naltrexone (1 mg/kg) and buprenorphine/naltrexone (1mg/kg + 1mg/kg). On day five, two hours after the last dose of the pretreatment drug, the dose-response curve for morphine was determined. Naltrexone shifted the morphine dose-response curve substantially to the right indicating blockade of opiate receptors whereas a combination of naltrexone (1 mg/kg) and buprenorphine (1 mg/kg) not only shifted the curve further to the right but also reduced the peak effect produced by morphine.

It is preferable to formulate the compositions in unit dosage forms ie physically discrete units containing the appropriate amounts of buprenorphine and naltrexone together with pharmaceutically acceptable diluents and/or carriers. Such compositions may be in the form of solid or liquid formulations.

Liquid preparations may be for example comprise buprenorphine or a non-toxic salt thereof plus naltrexone or a non-toxic salt thereof dissolved in 20-30% v/v aqueous ethanol buffered to between pH 4.5 to 5.5.

Compositions in the form of sublingual tablets contain soluble excipients such as lactose, mannitol, dextrose, sucrose or mixtures thereof. They will preferably also contain granulating and disintegrating agents such as starch, binding agents such as povidone or hydroxypropyl-methyl cellulose and lubricating agents such as magnesium stearate.

The invention is illustrated by the following Examples:-

EXAMPLE 1

10 ml of a sublingual solution containing 10 mg/ml buprenorphine and 10 mg/ml naltrexone in a pH5 mixture of a 30% v/v aqueous ethanol: citric acid/disodium hydrogen phosphate buffer was prepared as follows:

1. The buffer was prepared by mixing 3.8 ml 0.1 M citric acid and 3.2 ml 0.2 M disodium hydrogen phosphate.
2. 3.0 ml 95% v/v ethanol was added to the buffer increasing the pH from 4.6 to 5.0.
3. 108 mg buprenorphine hydrochloride was added with stirring until dissolved.

EP 0 319 243 B1

4. 110.7 mg naltrexone hydrochloride was added with stirring until dissolved.

5. Unit dose packs containing 0.2 ml were dispensed to give single doses of 2 mg buprenorphine and 2 mg naltrexone.

5 Unit dose packs containing 0.8 ml were dispensed to give single doses of 8 mg buprenorphine and 8 mg naltrexone.

EXAMPLE 2

10 A sublingual tablet having the following composition:

	mg/tablet
15 Buprenorphine HCl	2.16
Naltrexone HCl	2.21
20 Lactose	26.98
Mannitol	18.0
25 Maize starch	9.0
Povidone	1.2
30 Magnesium stearate	<u>0.45</u>
	60.0

35 was prepared by screening all the materials with the exception of the magnesium stearate through a 750µm sieve and blending them together. The mixed powders were then subjected to an aqueous granulation process and dried at 50°C. The resulting granules were forced through a 750µm sieve and blended with magnesium stearate (pre-sieved through a 500µm sieve). The tablet granules were compressed to yield tablets of 5.56 mm diameter and weight 60 mg.

40 EXAMPLE 3

The formulation of Example 2 was varied as follows, the method of manufacture being as for Example 2:

	mg/tablet
45 Buprenorphine HCl	8.64
50 Naltrexone HCl	8.84
Lactose	45.22
55 Mannitol	36.0
Maize starch	18.0
Povidone	2.4
60 Magnesium stearate	<u>0.90</u>
65	120.0

In this Example the tablet granules were compressed to yield tablets of 7 mm diameter and weight 120 mg.

EXAMPLE 4

The formulation of Example 3 was varied as follows. The method of manufacture being as for Example 2.

	mg/tablet
Buprenorphine HCl	4.32
Naltrexone HCl	4.42
Lactose	53.96
Mannitol	36.0
Maize starch	18.0
Povidone	2.4
Magnesium stearate	<u>0.90</u>
	120.0

In this Example the tablet granules were compressed to yield tablets of 7 mm diameter and weight 120 mg.

Claims

1. A pharmaceutical composition in sublingual unit dosage form for maintenance treatment of opiate addicts comprising from 2 to 8 mg buprenorphine and an amount of naltrexone sufficient to substantially attenuate the euphorogenic effect of the buprenorphine when injected and to provide greater opiate blocking effect than that of naltrexone alone wherein the weights of naltrexone to buprenorphine are within the ratio of 1:4 to 1:1, the weights of buprenorphine and naltrexone being expressed in terms of weights of the base forms.
2. A pharmaceutical composition in sublingual unit dosage form for maintenance treatment of opiate addicts comprising from 2 to 8 mg buprenorphine and an amount of naltrexone sufficient to substantially attenuate the euphorogenic effect of the buprenorphine when injected and to provide greater opiate blocking effect than that of naltrexone alone wherein the amount of naltrexone is in the range of 2 to 8 mg, the weights of buprenorphine and naltrexone being expressed in terms of weights of the base forms.
3. A pharmaceutical composition according to Claim 1 wherein the weight of buprenorphine is 4mg and that of naltrexone is in the range of 1 to 4 mg.
4. A pharmaceutical composition according to Claim 1 or Claim 2 wherein the weight of buprenorphine is 4 mg and that of naltrexone is in the range of 2 to 4 mg.

Patentansprüche

1. Pharmazeutische Zusammensetzung in sublingualer Einheitsdosisform zur Substitutionsbehandlung von Opiatsüchtigen, umfassend von 2 bis 8 mg Buprenorphin und eine Menge von Naltrexon, die ausreicht, um die euphorisierende Wirkung von injiziertem Buprenorphin wesentlich zu dämpfen und eine größere Opiat-blockierende Wirkung als diejenige von Naltrexon allein zu ergeben, worin die Gewichte von Naltrexon zu Buprenorphin im Verhältnis 1:4 bis 1:1 liegen und die Gewichte von Buprenorphin und Naltrexon als die Gewichte der Basenformen ausgedruckt sind.
2. Pharmazeutische Zusammensetzung in sublingualer Einheitsdosisform zur Substitutionsbehandlung von Opiatsüchtigen, umfassend von 2 bis 8 mg Buprenorphin und eine Menge von Naltrexon, die ausreicht, um die euphorisierende Wirkung von injiziertem Buprenorphin wesentlich zu dämpfen und eine größere Opiat-blockierende Wirkung als diejenige von Naltrexon allein zu ergeben, worin die Menge von

Naltrexon im Bereich von 2 bis 8 mg liegt und die Gewichte von Buprenorphin und Naltrexon als Gewichte der Basenformen ausgedrückt sind.

3. Pharmazeutische Zusammensetzung nach Anspruch 1, worin das Gewicht von Buprenorphin 4 mg beträgt und jenes von Naltrexon im Bereich von 1 bis 4 mg liegt.

5 4. Pharmazeutische Zusammensetzung nach Anspruch 1 oder 2, worin das Gewicht von Buprenorphin 4 mg beträgt und jenes von Naltrexon im Bereich von 2 bis 4 mg liegt.

Revendications

10 1. Composition pharmaceutique sous la forme d'une dose unitaire sublinguale pour le traitement d'entretien d'opiomanes comprenant de 2 à 8 mg de buprénorphine et une proportion de naltrexone qui suffit à sensiblement atténuer l'effet euphorigène de la buprénorphine lorsqu'elle est injectée et à assurer un effet de blocage des substances opiacées supérieur à celui de la naltrexone seule, caractérisée en ce que les poids de la naltrexone à la buprénorphine se situent dans le rapport de 1:4 à 1:1, les poids de la buprénorphine et de la naltrexone s'exprimant en poids des formes base.

15 2. Composition pharmaceutique sous la forme d'une dose unitaire sublinguale pour le traitement d'entretien d'opiomanes comprenant de 2 à 8 mg de buprénorphine et une proportion de naltrexone qui suffit à sensiblement atténuer l'effet euphorigène de la buprénorphine lorsqu'elle est injectée et à assurer un effet de blocage des substances opiacées supérieur à celui de la naltrexone seule, caractérisée en ce que la proportion de la naltrexone varie de 2 à 8 mg, les poids de la buprenorphine et de la naltrexone s'exprimant en poids des formes base.

20 3. Composition pharmaceutique suivant la revendication 1, caractérisée en ce que le poids de la buprénorphine est de 4 mg et en ce que celui de la naltrexone fluctue de 1 à 4 mg.

25 4. Composition pharmaceutique suivant la revendication 1 ou la revendication 2, caractérisée en ce que le poids de la buprénorphine est de 4 mg et en ce que celui de la naltrexone fluctue de 2 à 4 mg.

30

35

40

45

50

55

60

65